



HANDBOOK



No. 12-10

Mar 12



Tactical Combat Casualty Care

Observations, Insights, and Lessons

U.S. UNCLASSIFIED
REL NATO, GCTF, ISAF, ABCA
FOR OFFICIAL USE ONLY

Handling Instructions for CALL Electronic Media and Paper Products

Center for Army Lessons Learned (CALL) authorizes official use of this CALL product for operational and institutional purposes that contribute to the overall success of U.S., coalition, and allied efforts.

The information contained in this product reflects the actions of units in the field and may not necessarily be approved U.S. Army policy or doctrine.

This product is designed for official use by U.S., coalition, and allied personnel and cannot be released to the public without the consent of CALL. This product has been furnished with the expressed understanding that it will be used for official defense-related purposes only and that it will be afforded the same degree of protection that the U.S. affords information marked "U.S. UNCLASSIFIED, For Official Use Only [FOUO]" in accordance with U.S. Army Regulation (AR) 380-5, section 5-2.

Official military and civil service/government personnel, to include all coalition and allied partners, may paraphrase; quote; or use sentences, phrases, and paragraphs for integration into official products or research. However, integration of CALL "U.S. UNCLASSIFIED, For Official Use Only [FOUO]" information into official products or research renders them FOUO, and they must be maintained and controlled within official channels and cannot be released to the public without the consent of CALL.

This product may be placed on protected UNCLASSIFIED intranets within military organizations or units, provided that access is restricted through user ID and password or other authentication means to ensure that only properly accredited military and government officials have access to these products.

Regulations strictly forbid posting CALL "U.S. UNCLASSIFIED, For Official Use Only [FOUO]" documents to Department of Defense (DOD) websites that do not restrict access to authorized personnel. AR-25-1, 15 Jul 2005, Army Knowledge Management and Information Technology, paragraph 6-4 n (2) (b) and DOD Web Site Administration Policy and Procedures (11 Jan 2002), Part II, paragraph 3.6.1 require appropriate mechanisms to protect sensitive information.

When no longer needed, all CALL "U.S. UNCLASSIFIED, For Official Use Only [FOUO]" paper products and electronic media will be shredded or destroyed using approved paper shredders or CDROM destroyers.

To allied and coalition personnel:

This information is furnished with the understanding that it is to be used for defense purposes only, that it is to be afforded essentially the same degree of security protection as such information is afforded by the United States, and that it is not to be revealed to another country or international organization without the consent of CALL.

CENTER FOR ARMY LESSONS LEARNED

SUPPORTING THE WARFIGHTER



Tactical Combat Casualty Care Handbook

DIGITAL VERSION AVAILABLE

A digital version of this CALL publication is available to view, download, or reproduce from the CALL restricted website, <<http://call.army.mil>>. Reproduction of this publication is welcomed and highly encouraged.

Common Access Card (CAC) or Army Knowledge Online (AKO) login is required to access the digital version.

This publication is located online at:

**[https://call2.army.mil/toc.aspx?document=6851&
filename=/docs/doc6851/12-10.pdf](https://call2.army.mil/toc.aspx?document=6851&filename=/docs/doc6851/12-10.pdf)**

U.S. UNCLASSIFIED
REL NATO, GCTF, ISAF, ABCA
For Official Use Only



Foreword

A decade of intense combat in two theaters has taught us many lessons about what works and what does not in the effort to accomplish that all-important mission of saving lives in battle. A severely injured Soldier today has about twice the likelihood of surviving his wounds compared to Soldiers in wars as recent as Vietnam. That progress is the result of many things: better tactics and weapons, better body armor and helmets, better trained and fitter Soldiers. But, the introduction of tactical combat casualty care (TCCC) throughout the Army has certainly been an important part of that improvement.

TCCC is fundamentally different from civilian care. It is the thoughtful integration of tactics and medicine, but to make it work takes a different set of skills and equipment, and every Soldier and leader needs to understand it and practice it.

This handbook is the result of years of careful study of the care of wounded Soldiers, painstaking research by medics and physicians, and the ability of leaders at all levels to see and understand the lessons being learned and the willingness to make the changes in equipment, training, and doctrine needed to improve the performance of the Army Health System. It is the best guidance we have at the time of publication, but new information, new techniques, or new equipment will drive changes in the future. Be assured that these performance improvement efforts will continue as long as American Soldiers go in harm's way.

JAMES W. KIRKPATRICK, MD
COL (ret), MC
Senior Clinical Consultant
Directorate of Combat and Doctrine Development
AMEDD Center & School

Tactical Combat Casualty Care Handbook	
Table of Contents	
Chapter 1. Tactical Combat Casualty Care	1
Chapter 2. Tactical Combat Casualty Care Procedures	35
Appendix A. Triage Categories	57
Appendix B. Evacuation Categories	59
Appendix C. 9-Line Medical Evacuation	61
Appendix D. Combat Pill Pack	63
Appendix E. Improved First-Aid Kit	65
Appendix F. Warrior Aid and Litter Kit	67
Appendix G. Aid Bag Considerations	69
Appendix H. National Stock Numbers	71
Appendix I. References/Resources	73

Center for Army Lessons Learned	
Director	COL Thomas H. Roe
Sustainment WfF Team Chief	LTC Vaughn M. Grizzle
CALL Analysts	Donald Haus Jose Jimenez, CALL Contractor
Contributing Authors	Donald L. Parsons, PA-C, MPAS Jeffrey Mott, PA-C, MPAS
Editor	Jenny Solon
Graphic Artist	Eric Eck, CALL Contractor

CALL wishes to acknowledge the Army Medical Department Center and School – Lessons Learned Division, the Committee on Tactical Combat Casualty Care, and the Army Institute of Surgical Research for their assistance in compiling this handbook.

CENTER FOR ARMY LESSONS LEARNED

The Secretary of the Army has determined that the publication of this periodical is necessary in the transaction of the public business as required by law of the Department.

Unless otherwise stated, whenever the masculine or feminine gender is used, both are intended.

Note: Any publications (other than CALL publications) referenced in this product, such as ARs, FMs, and TMs, must be obtained through your pinpoint distribution system.

Chapter 1

Tactical Combat Casualty Care

Section I: Introduction

Tactical combat casualty care (TCCC) is the pre-hospital care rendered to a casualty in a tactical, combat environment. The principles of TCCC are fundamentally different from those of traditional civilian trauma care where most medical providers and medics train. These differences are based on both the unique patterns and types of wounds that are suffered in combat and the tactical conditions medical personnel face in combat. Unique combat wounds and tactical conditions make it difficult to determine which intervention to perform at what time. Besides addressing a casualty's medical condition, responding medical personnel must also address the tactical situation faced while providing casualty care in combat. A medically correct intervention performed at the wrong time may lead to further casualties. Put another way, "good medicine may be bad tactics," which can get the rescuer and casualty killed. To successfully navigate these issues, medical providers must have skills and training oriented to combat trauma care, as opposed to civilian trauma care.

Casualties and Wounds

On the battlefield, the pre-hospital period is the most important time to care for any combat casualty. In previous wars, up to 90 percent of combat deaths occurred before a casualty reached a medical treatment facility. This highlights the primary importance of treating battlefield casualties at the point of injury, prior to casualty evacuation and arrival at a treatment facility.

Specifically, combat deaths result from the following:

- 31 percent: Penetrating head trauma.
- 25 percent: Surgically uncorrectable torso trauma.
- 10 percent: Potentially correctable surgical trauma.
- **9 percent: Exsanguination.**
- 7 percent: Mutilating blast trauma.
- **3-4 percent: Tension pneumothorax (PTX).**
- **2 percent: Airway obstruction/injury.**
- 5 percent: Died of wounds (mainly infection and shock).

(**Note:** Numbers do not add up to 100 percent. Not all causes of death are listed. Some deaths are due to multiple causes.)

A significant percentage of these deaths (highlighted above in bold type) are potentially avoidable with proper, timely intervention. Of these avoidable deaths, the vast majority are due to exsanguination and airway or breathing difficulties, conditions that can and should be addressed at the point of injury. It has been estimated that of all preventable deaths, up to 90 percent of them can be avoided with the simple application of a tourniquet for extremity hemorrhage, the rapid treatment of a PTX, and the establishment of a stable airway.

On the battlefield, casualties will fall into three general categories:

- Casualties who will die, regardless of receiving any medical aid.
- Casualties who will live, regardless of receiving any medical aid.
- Casualties who will die if they do not receive timely and appropriate medical aid.

TCCC addresses the third category of casualties — those who require the most attention of the medical provider during combat.

TCCC versus Advanced Trauma Life Support

Trauma care training for military medical personnel traditionally has been based on the principles of the civilian Emergency Medical Technicians Basic Course and basic and advanced trauma life support (ATLS). These principles, especially ATLS, provide a standardized and very successful approach to the management of civilian trauma patients in a hospital setting. However, some of these principles may not apply in the civilian pre-hospital setting, let alone in a tactical, combat environment.

The pre-hospital phase of casualty care is the most critical phase of care for combat casualties, accounting for up to 90 percent of combat deaths. Furthermore, combat casualties can suffer from potentially devastating injuries not usually seen in the civilian setting. Most casualties during combat are the result of penetrating injuries, rather than the blunt trauma seen in the civilian setting. Combat casualties may also suffer massive, complex trauma, such as traumatic limb amputation. In addition to the medical differences between civilian and combat trauma, several other factors affect casualty care in combat, including the following:

- Hostile fire may be present, preventing the treatment of the casualty.
- Medical equipment is limited to that carried by mission personnel.

- Tactical considerations may dictate that mission completion take precedence over casualty care.
- Time until evacuation is highly variable (from minutes to hours or days).
- Rapid evacuation may not be possible based on the tactical situation.

TCCC Goals

TCCC presents a system to manage combat casualties that considers the issues discussed above. An important guiding principle of TCCC is performing the correct intervention at the correct time in the continuum of field care. To this end, TCCC is structured to meet three important goals:

- Treat the casualty.
- Prevent additional casualties.
- Complete the mission.

Stages of Care

In thinking about the management of combat casualties, it is helpful to divide care into three distinct phases, each with its own characteristics and limitations:

- **Care under fire** is the care rendered at the point of injury while both the medic and the casualty are under effective hostile fire. The risk of additional injuries from hostile fire at any moment is extremely high for both the casualty and the medic. Available medical equipment is limited to that carried by the medic and the casualty.
- **Tactical field care** is the care rendered by the medic once he and the casualty are no longer under effective hostile fire. It also applies to situations in which an injury has occurred on a mission but there has been no hostile fire. Available medical equipment is still limited to that carried into the field by mission personnel. Time to evacuation may vary from minutes to hours.
- **Tactical evacuation care** is the care rendered once the casualty has been picked up by an aircraft, vehicle, or boat. Additional medical personnel and equipment that has been pre-staged in these assets should be available during this phase of casualty management.

The chapters and sections of this handbook will present a discussion of each stage of TCCC as well as instructions for the procedures TCCC requires.

Section II: Care Under Fire

Care under fire is the care rendered by the rescuer at the point of injury while he and the casualty are still under effective hostile fire. The risk of additional injuries at any moment is extremely high for both the casualty and the rescuer. The major considerations during this phase of care are the following:

- Suppression of hostile fire.
- Moving the casualty to a safe position.
- Treatment of immediate life-threatening hemorrhage.

Casualty care during the care under fire phase is complicated by several tactical factors. First, the medical equipment available for care is limited to that which is carried by the individual Soldiers and the rescuers. Second, the unit's personnel will be engaged with hostile forces and, especially in small-unit engagements, will not be available to assist with casualty treatment and evacuation. Third, the tactical situation prevents the medic or medical provider from performing a detailed examination or definitive treatment of casualties. Furthermore, these situations often occur during night operations, resulting in severe visual limitations while treating the casualty.

Defensive Actions

The best medicine on the battlefield is fire superiority. The rapid success of the combat mission is the immediate priority and the best way to prevent the risk of injury to other personnel or additional injuries to casualties. Medical personnel carry small arms to defend themselves and casualties in the field. The additional firepower from the medical personnel may be essential to obtaining fire superiority. Initially, medical personnel may need to assist in returning fire before stopping to care for the casualty. Additionally, casualties who have sustained non-life-threatening injuries and are still able to participate in the fight must continue to return fire as they are able.

As soon as the rescuer is directed or able, his first major objective is to keep the casualty from sustaining additional injuries. Wounded Soldiers who are unable to participate further in the engagement should lay flat and still if no ground cover is available or move as quickly as possible if nearby cover is available. The medic may be able to direct the injured Soldier to provide self-aid.

Airway Management

Do not perform any immediate management of the airway during the care under fire phase. Airway injuries typically play a minimal role in combat casualties, comprising only 1 to 2 percent of casualties primarily from

maxillofacial injuries. The primary concern is to move the casualty to cover as quickly as possible. The time, equipment, and positioning required to manage an impaired airway expose the casualty and rescuer to increased risk. Rescuers should defer airway management until the tactical field care phase, when the casualty and rescuer are safe from hostile fire.

Hemorrhage Control

The number one cause of preventable battlefield deaths is hemorrhage from compressible wounds. Therefore, the primary medical interventions during the care under fire phase are directed toward stopping any life-threatening bleeding as quickly as possible. Injuries to an artery or other major vessel can rapidly result in hemorrhagic shock and exsanguinations. A casualty may exsanguinate before medical help arrives, so definitive control of life-threatening hemorrhage on the battlefield cannot be overemphasized. In Vietnam, bleeding from an extremity wound was the cause of death in more than 2,500 casualties who had sustained no other injury.

Extremity wounds. The rapid, temporary use of a tourniquet is the recommended management for all life-threatening extremity hemorrhage. Standard field dressings and direct pressure may not work reliably to control extremity hemorrhage. While traditional ATLS training discourages the use of tourniquets, they are appropriate in the tactical combat setting. The benefits of tourniquet use over other methods of hemorrhage control include:

- Direct pressure and compression are difficult to perform and maintain in combat settings and result in delays in getting the rescuer and casualty to cover.
- Tourniquets can be applied to the casualty by himself, thus limiting the rescuer's exposure to hostile fire.
- There are few complications from tourniquet use. Ischemic damage is rare if the tourniquet is in place for less than two hours.

During the care under fire phase, the casualty and rescuer remain in grave danger from hostile fire. If the casualty is observed to have bleeding from an extremity, the care provider should apply a tourniquet to the injured extremity over the uniform, high on the extremity, and move himself and the casualty to cover as quickly as possible.

Nonextremity wounds. These injuries are difficult to treat in the care under fire phase. Attempt to provide direct pressure to these wounds as you rapidly move the casualty to cover. Once under cover, a hemostatic agent is appropriate for these injuries.

Casualty Transportation

Transportation of the casualty is often the most problematic aspect of TCCC. In the care under fire phase, transportation is complicated by the limited equipment and personnel available and the risk of further injury due to hostile fire. Removing the casualty from the field of fire as quickly as possible is the transportation priority during this phase of care. Do not attempt to save a casualty's rucksack unless it contains items that are critical to the mission. However, if at all possible, take the casualty's weapons and ammunition. The enemy may use them against you.

Cervical spine immobilization. Although the civilian standard of care is to immobilize the cervical spinal column prior to moving a patient with injuries that might have resulted in damage to the spine, this practice is generally not appropriate in the combat setting. In Vietnam, studies examining the value of cervical spinal immobilization in penetrating neck injuries found that only 1.4 percent of casualties with penetrating neck injuries would have possibly benefited from immobilization of the cervical spine. The time required to accomplish cervical spine immobilization was found to be 5.5 minutes, even when using experienced rescuers. In addition, the equipment needed for this procedure (long spine board) is generally not available at the point of wounding. Therefore, the potential hazards of hostile fire to both the casualty and rescuer outweigh the potential benefit of cervical spine immobilization. However, for casualties with significant blunt trauma, cervical spine immobilization is appropriate during the care under fire phase. Parachuting injuries, fast-roping injuries, falls greater than 15 feet, and other types of trauma resulting in neck pain or unconsciousness should be treated with spinal immobilization, unless the danger of hostile fire constitutes a greater risk in the judgment of the medic.

Transportation methods. Standard litters for patient evacuation may not be available for movement of casualties in the care under fire phase. Consider using alternate methods of evacuation, such as Sked or Talon II litters or dragging the casualty out of the field of fire by his web gear, poncho, or even a length of rope with a snap link. There are a number of drag straps and drag litters available to help expedite this move. Traditional one- and two-man carries are not recommended, as the weight of the average combatant makes these types of casualty movement techniques extremely difficult. Additionally, consider the use of obscurants such as smoke or CS (irritating agent) to assist in casualty recovery. Vehicles can also be used as a screen during recovery attempts. In Iraq, there have been several instances of tanks being used as screens to facilitate casualty evacuation (CASEVAC).

Section III: Tactical Field Care

Tactical field care is the care rendered to the casualty once the casualty and rescuer are no longer under effective hostile fire. This term also applies to situations in which an injury has occurred on a mission but there has been no hostile fire. This phase of care is characterized by the following:

- The risk from hostile fire has been reduced but still exists.
- The medical equipment available is still limited by what has been brought into the field by mission personnel.
- The time available for treatment is highly variable. Time prior to evacuation, or re-engagement with hostile forces, can range from a few minutes to many hours.

Medical care during this phase of care is directed toward more in-depth evaluation and treatment of the casualty, focusing on those conditions not addressed during the care under fire phase of treatment. While the casualty and rescuer are now in a somewhat less hazardous situation, this is still not the setting for a true rapid trauma assessment and treatment. Evaluation and treatment are still dictated by the tactical situation.

In some cases, tactical field care will consist of rapid treatment of wounds with the expectation of a re-engagement with hostile forces at any moment. The need to avoid undertaking nonessential evaluation and treatment is critical in such cases. Conversely, care may be rendered once the mission has reached an anticipated evacuation point without pursuit and is awaiting evacuation. In these circumstances, there may be ample time to render whatever care is feasible in the field. However, as time to evacuation may vary greatly, medical providers and medics must take care to partition supplies and equipment in the event of prolonged evacuation wait times.

CPR

In casualties of blast or penetrating injury found to be without pulse, respiration, or other signs of life, CPR on the battlefield will generally not be successful and should not be attempted. Attempts to resuscitate trauma patients in arrest have been found to be futile even in urban settings where victims are in close proximity to trauma centers. On the battlefield, the cost of performing CPR on casualties with what are inevitably fatal injuries will result in additional lives lost as care is withheld from casualties with less severe injuries. Also, these attempts expose rescuers to additional hazards from hostile fire. Prior to the tactical evacuation care phase, rescuers should consider CPR only in the cases of nontraumatic disorders such as hypothermia, near drowning, or electrocution.

Altered Mental Status

Immediately disarm any casualty with an altered mental status, including secondary weapons and explosive devices. An armed combatant with an altered mental status is a significant risk to himself and those in his unit. The four main reasons for an altered mental status are traumatic brain injury (TBI), pain, shock, and analgesic medication (for example, morphine).

Hemorrhage Control

In the tactical field care phase, hemorrhage control includes addressing any significant bleeding sites not previously controlled. When evaluating the casualty for bleeding sites, only remove the absolute minimum of clothing needed to expose and treat injuries. Stop significant extremity bleeding as quickly as possible, using a tourniquet without hesitation if necessary. It is important to note that after tourniquet application, a distal pulse must be assessed to ensure the arterial blood flow has been stopped. If a distal pulse remains after tourniquet application, then a second tourniquet must be applied side by side and just above the original tourniquet. This second tourniquet applies pressure over a wider area and more easily stops the arterial flow.

There have been a number of reports of compartment syndrome in distal extremities when the tourniquet is not applied tightly enough to stop arterial blood flow. In addition, there have been tourniquet failures when the care provider has attempted to tighten the tourniquet to the extreme. If a tourniquet is applied around the limb as snugly as possible before the windlass is tightened, it should only take three revolutions (540 degrees) of the windlass to stop blood flow. If a distal pulse is still present, it is more prudent to apply a second tourniquet as described above than to try and tighten the original one too tightly. It must be pointed out that the additional step of checking a distal pulse should only be accomplished when the tactical situation permits. Otherwise, direct pressure, pressure dressings, or homeostatic dressings (combat gauze) should be used to control bleeding.

Tourniquets should remain in place until the casualty has been transported to the evacuation point. Once the patient has been transported to the site where evacuation is anticipated, and any time the casualty is moved, reassess any tourniquets previously applied. If evacuation is significantly delayed (greater than two hours), the medic should make a determination if the tourniquet should be loosened and bleeding control replaced with some other technique. Hemostatic bandages, pressure bandages, etc. may be able to control the bleeding and lower the risk to the extremity that a tourniquet poses. However, it needs to be emphasized that there is no evidence that tourniquets have caused the loss of any limbs in hundreds of tourniquet applications. If a decision to remove a tourniquet is made, the medic must be sure to complete any required fluid resuscitation prior to tourniquet

discontinuation. It is not necessary to remove the tourniquet, only to loosen it. This allows the tourniquet to be reapplied if the hemorrhage cannot be controlled by other methods.

Data from research done in theater have demonstrated tourniquet application before the casualty goes into shock significantly improves survival statistics. The training emphasis must continue to be on the control of bleeding in all casualties.

Airway Management

In the tactical field care phase, direct initial management to the evaluation and treatment of the casualty's airway once all hemorrhage problems have been addressed. Intervention should proceed from the least invasive procedure to the most invasive. Do not attempt any airway intervention if the casualty is conscious and breathing well on his own. Allow the casualty to assume the most comfortable position that best protects his airway, to include sitting upright.

Unconscious casualty without airway obstruction. If the casualty is unconscious, the most likely cause is either hemorrhagic shock or head trauma. In either case, an adequate airway must be maintained. If the unconscious casualty does not exhibit signs of airway obstruction, the airway should first be opened with a chin lift or a jaw-thrust maneuver. As in the care under fire phase, cervical spine immobilization is generally not required, except in the instance of significant blunt trauma.

If spontaneous respirations are present without respiratory distress, an adequate airway in the unconscious casualty is best maintained with a nasopharyngeal airway (NPA). An NPA is preferred over an oropharyngeal airway because it is better tolerated if the casualty regains consciousness and is less likely to be dislodged during casualty transport. After inserting the NPA, place the casualty in the recovery position (see Figure 1-1) to maintain the open airway and prevent aspiration of blood, mucous, or vomit.



Figure 1-1. Recovery position

Current or impending airway obstruction. For casualties with a current or impending airway obstruction, the initial intervention is again to open the airway with either a chin lift or a jaw-thrust maneuver. Either maneuver

is followed by the insertion of an NPA. However, if an airway obstruction develops or persists despite the use of an NPA, a more definitive airway is required. In some casualties a more definitive airway may consist of a supraglottic device, such as a combitube or King LT. These airways are not well tolerated unless the casualty is totally obtunded. These devices are easily inserted and able to maintain an open airway better than a simple NPA. However, often a surgical cricothyroidotomy may be indicated.

Cricothyroidotomy. Significant airway obstruction in the combat setting is likely the result of penetrating wounds of the face or neck, where blood or disrupted anatomy precludes good visualization of the vocal cords. This setting makes endotracheal intubation highly difficult, if not impossible. In these cases, surgical cricothyroidotomy is preferable over endotracheal intubation. This procedure has been reported safe and effective in trauma victims, and in the hands of a rescuer who does not intubate on a regular basis, it should be the next step when other airway devices are not effective. Furthermore, cricothyroidotomy can be performed under local anesthesia with lidocaine on a casualty who is awake. The majority of preventable airway deaths occurred from penetrating trauma to the face and neck, where disrupted anatomy and significant bleeding made airway interventions very difficult.

Intubation. Endotracheal intubation is the preferred airway technique in civilian trauma settings, but this procedure may be prohibitively difficult in the tactical environment. Many medics have never intubated a live person; their experience is only with mannequins in a controlled environment and is infrequent at best. The standard endotracheal intubation technique requires the use of tactically compromising white light. Also, esophageal intubations are more likely with the inexperienced intubator and much more difficult to detect in the tactical environment. Finally, most airway obstructions on the battlefield are the result of penetrating wounds of the head and neck, where cricothyroidotomy is the procedure of choice.

Breathing

The next aspect of casualty care in the tactical field care phase is the treatment of any breathing problems, specifically the development of either an open PTX or a tension PTX.

Penetrating chest wounds. Traumatic defects in the casualty's chest wall may result in an open PTX. All open chest wounds should be treated as such. Cover the wound during expiration with an occlusive dressing; numerous different materials are available for use. In addition, multiple commercial chest seals are now available, many with excellent adhesive properties. The dressing should be sealed on all four sides. The casualty should then be placed in a sitting position, if applicable, and monitored for

the development of a tension PTX, which should be treated as described below.

Tension PTX. Assume any progressive, severe respiratory distress on the battlefield resulting from unilateral penetrating chest trauma represents a tension PTX. Do not rely on such typical signs as breath sounds, tracheal shift, and hyper-resonance on percussion for diagnosis in this setting, because these signs may not always be present. Even if these signs are present, they may be difficult to detect on the battlefield. Treat tension PTXs in the tactical field care phase via decompression with a 14-gauge, 3.25-inch-long needle catheter. A casualty with penetrating chest trauma will generally have some degree of hemothorax or PTX as a result of his primary wound. The additional trauma caused by a needle thoracostomy will not worsen his condition should he not have a tension PTX. Decompress the casualty with the needle and catheter, removing the needle and leaving the catheter buried to the hub. The medic must monitor this casualty after the procedure to ensure the catheter has not clotted or dislodged and that respiratory symptoms have not returned. If respiratory symptoms have returned or the catheter is clotted or dislodged, flush the catheter or perform a second needle thoracostomy adjacent to the first. Chest tubes are not recommended during this phase of care, as they are not needed for initial treatment of a tension PTX, are more technically difficult and time-consuming to perform, and are more likely to result in additional tissue damage and subsequent infection.

Vascular Access

Obtain IV access at this point during the tactical field care phase. While ATLS training teaches starting two large-bore (14- or 16-gauge) IV catheters, the use of a single 18-gauge catheter is preferred in the tactical setting. The 18-gauge catheter is adequate for rapid delivery of resuscitation fluids and medication, is easier to insert, and conserves the supplies in a medic aid bag. Medics should not start an IV on an extremity that may have a significant wound proximal to the IV insertion site.

If the casualty requires fluid resuscitation and IV access cannot be obtained, sternal intraosseous (IO) access is recommended. One possible IO fluid delivery system is the First Access for Shock and Trauma (FAST1) System. Other extremity IO devices are available, but it should be remembered that the majority of injuries are penetrating lower extremity injuries. Cutdowns are not recommended in the tactical setting as they are time-consuming, technically difficult, and require instruments that in all likelihood will not be available. Medics will most likely not be trained, equipped, or authorized to perform cutdowns.

Fluid Resuscitation

Fluid resuscitation during the tactical field care phase is significantly different than in the civilian pre-hospital setting. During this phase of care, fluid resuscitation is guided by several assumptions:

- The tactical situation may not allow time for thorough fluid resuscitation. Care may consist only of immediate evacuation while in extremis.
- Lack of hemorrhage control is the leading cause of preventable death on the battlefield. Therefore, hemorrhage control is paramount and takes priority over fluid resuscitation, especially in a situation with limited time and resources.
- Stethoscopes, blood pressure cuffs, and other equipment used in the hospital setting to monitor fluid status and shock are rarely available or useful in a noisy and chaotic battlefield setting. In the tactical setting, assessing a casualty's mental status and peripheral pulses is adequate for determining the need for fluid resuscitation.

In light of these considerations, during the tactical field care phase, only provide fluid resuscitate to those casualties exhibiting signs of shock or TBI. If the casualty has only superficial wounds, IV fluid resuscitation is not necessary, but oral fluids should be encouraged. In those casualties with significant wounds who are coherent and without any obvious blood loss or signs of shock, blood loss likely has been stopped. In these casualties, obtain IV and/or IO access, hold IV fluids, and re-evaluate the casualty as frequently as possible.

Shock. Shock encountered in the combat setting will most likely be hemorrhagic shock. Assume the casualty is in shock if he has an altered mental status in the absence of head injury and/or has weak or absent peripheral pulses. Begin fluid resuscitation in these cases.

- During the tactical field care phase, 6 percent Hetastarch (Hextend) is the recommended fluid for resuscitation. Hextend is preferred over crystalloid fluids because one 500-milliliter (ml) bag of Hextend is physiologically equivalent to three 1,000-ml bags of lactated Ringer's solution (LRS), weighs 5.5 pounds less, and expands intravascular fluid volume for at least 8 hours.
- Initiate fluid resuscitation with a 500-ml Hextend IV bolus.
- Monitor the casualty, and, if after 30 minutes the casualty still has no peripheral pulse or has altered mentation, administer a second 500-ml Hextend bolus.

- Do not administer more than 1,000 ml of Hextend. This is equivalent to six liters of LRS.

If the casualty is still in shock after 1,000 ml of Hextend, the casualty is probably still bleeding. Fluid resuscitation is unlikely to be effective until the hemorrhage is controlled. The casualty needs to be evacuated to surgical care as soon as possible. If rapid evacuation is not feasible, the medic may need to consider triaging medical supplies and focusing attention on more salvageable casualties.

TBI. Head injuries are special situations. Hypotension and hypoxia exacerbate secondary brain injury and are difficult to control in the initial phases of combat casualty care. If the casualty with a TBI is unconscious and has no peripheral pulse, he should be resuscitated to restore a palpable radial pulse and evacuated as soon as possible.

Hypothermia Prevention

Combat casualties are at a high risk for hypothermia, which is defined as a whole body temperature below 95 F (35 C). Hypothermia can occur regardless of the ambient temperature. The blood loss typically associated with combat trauma results in peripheral vasoconstriction, which contributes to the development of hypothermia. In addition, the longer a casualty is exposed to the environment during treatment and evacuation, especially in wet conditions, the more likely the development of hypothermia. This is even more the case during rotary-wing evacuation.

Hypothermia, acidosis, and coagulopathy constitute the “lethal triad” in trauma patients. The association of hypothermic coagulopathy with high mortality has been well described. Hypothermia causes the inhibition of coagulation proteins, thus exacerbating the bleeding problem. The need to prevent hypothermia is highlighted by the fact that up to 10 percent of combat casualties arrive at a level III treatment facility exhibiting some degree of hypothermia.

During the tactical field care phase, the rescuer must first minimize the casualty’s exposure to the elements. If possible, keep all protective gear on the patient. However, if at all possible, replace any wet clothing. Use any methods available to keep the casualty warm, such as dry blankets, poncho liners, and sleeping bags.

One product now available is the hypothermia prevention and management kit (HPMK), which should be carried by all medics. The HPMK consists of a Ready-Heat blanket, which actively warms to 110-118 F when opened, and a reinforced heat-reflective shell. When the casualty is ready for transport, place the Ready-Heat blanket around his torso, and then place the casualty into the heat-reflective shell.

Monitoring/Further Evaluation

During the tactical field care phase, monitor the casualty clinically, and frequently reassess him until evacuation. Pulse oximetry, at a minimum, should be included in the medic aid bag and used as an adjunct to clinical monitoring. Keep in mind, though, that pulse oximetry readings may be misleading in the settings of shock and hypothermia. With the advent of operations in areas of high altitude, care should prevail with the interpretation of pulse oximetry readings at extreme elevations.

Carefully check the casualty for additional wounds. High-velocity projectiles from assault rifles may tumble and take erratic courses in tissue, leading to exit sites removed from the entry wound. Inspect and dress all wounds.

Pain Control

All casualties in pain should be given analgesia. The type and route of medication is dependent upon whether the casualty is conscious, still able to fight, and if IV access has been obtained.

Able to fight. If the casualty is conscious and still able to fight, give him oral pain medications that will not alter his level of consciousness. The recommended medications are Meloxicam (Mobic), 15 milligrams (mg) once daily, along with two 650-mg bilayered Tylenol caplets. These medications, along with an oral antibiotic, make up the combat pill pack, which all combatants should be instructed to take when they sustain a penetrating wound on the battlefield.

Unable to fight. If the casualty is seriously injured, in pain, and otherwise unable to fight, he should be given narcotic medications. Medics must be trained in the use of Naloxone (Narcan) and have it readily available before administering any narcotics. Closely monitor the casualty for any respiratory depression. Clearly and visibly document the use of any narcotics to avoid overdose and respiratory compromise.

- IV/IO access present: Administer an initial dose of 5 mg morphine Sulfate IV. This can be repeated every 10 minutes as needed to control severe pain. Accompany morphine use with 25 mg of Promethazine IV/IO/intramuscular (IM) every 4 hours as needed to control nausea and vomiting.

Fractures

Splint all fractures as circumstances allow, ensuring that peripheral pulse, sensory, and motor checks are performed both before and after splinting.

Be aware of possible compartment syndrome with suspected fractures associated with blast-injured Soldiers. The absence of a distal pulse with a possible fracture should be cause for more immediate evacuation.

Infection Control

Infection is an important cause of morbidity and mortality in battlefield wounds. Assume all open wounds on the battlefield are infected, and treat them with antibiotics. Choose antibiotics that cover a broad spectrum of organisms, with the specific medications based on available delivery route and any medication allergies the casualty may have.

Able to take oral medication. If the casualty can take oral medications, 400 mg of Moxafloxacin, taken once daily, is recommended. This medication should be part of the combat pill pack. The casualty should take this as soon as he is injured and all life-threatening injuries have been addressed.

Unable to take oral medication. If the casualty is unable to take oral medications because of shock, unconsciousness, or other reasons, IV or IM antibiotics should be given. Recommended antibiotics in this case include Cefotetan, 2 grams (g) IV (over 3 to 5 minutes) or IM every 12 hours; or Ertapenem, 1 g IV (over 30 minutes) or IM every 24 hours.

Penetrating Eye Trauma

Penetrating eye trauma presents a problem with care providers on the battlefield. These injuries can deteriorate without proper care. If a penetrating eye injury is suspected, perform a rapid field test of the individual's visual acuity. It is not necessary to use a vision (Snellen) chart to do so. Have the patient read any printed material or try to determine how many fingers you are holding up, or see if the patient can distinguish between light and dark. If vision is impaired, apply an eye shield to the eye (not a pressure bandage). Avoid any pressure being placed on the eye, as this could cause the internal contents of the eye to be pushed out. If available, give the casualty a 400-mg Moxifloxacin tablet to provide antibiotic coverage.



Figure 1-2



Figure 1-3

Burns

Burn casualties should have their wounds covered with dry sterile bandages. Avoid using “WaterGel” directly on the burns. Calculate the total body surface area (TBSA) of the burn by using the “Rule of Nine.”

Burns to the face and neck should raise the suspicion for airway compromise, and the care provider should be prepared to initiate airway support if necessary.

Fluid resuscitation should be accomplished by using the new “Rule of Ten.” Fluid resuscitation should be necessary for TBSA burns greater than 20 percent.

Rule of Ten

Calculate the TBSA of the burns to the nearest ten percent.

- Example: 43% TBSA burn would become 40%, 46% TBSA burn would become 50%.

Fluid resuscitation protocol:

Initial IV/IO fluid rate is calculated as percent TBSA x 10cc/hour for adults weighing 40-80 kilograms (kg). For every 10 kg above 80 kg, increase initial rate by 100 ml/hour. Examples:

- Casualty who weighs 50 kg and has a 40% TBSA burn; $40 \times 10 \text{ ml} = 400 \text{ ml}$ per hour. If possible, monitor urine output to 30-50 ml/hour.
- If casualty is 90 kg with a 40% TBSA, the formula would be $40 \times 10 \text{ mls} = 400 \text{ ml/hour} + 100 \text{ ml}$, for a total of 500 ml/hour.

The fluid of choice for isolated burns is LRS. If the casualty has additional wounds and has lost blood, Hextend may be used to prevent or treat shock. The amount of Hextend used should not exceed 1,000 ml as in the shock protocol.

Analgesia for burns should follow the guidelines in the paragraphs above for significant pain.

Antibiotics are not required for burns alone, but may be appropriate for other penetrating injuries.

The key to successful burn management is to evacuate the casualty to definitive care as rapidly as possible.

Spinal Precautions

Care under fire. Direct the casualty to move to cover and apply self-aid if able. If the casualty requires assistance, move him to cover. If the mechanism of injury included blunt trauma (such as riding in a vehicle that was struck by an improvised explosive device), minimize spinal movement while extricating him from the vehicle and moving him to cover. The casualty should be moved along his long spinal axis if at all possible while attempting to stabilize the head and neck.

Tactical field care, tactical evacuation care. Use spinal motion restriction techniques as defined below for casualties whose mechanism of injury included blunt trauma if (a) they are unconscious; (b) they are conscious

and have midline cervical spine tenderness or midline back pain; or (c) they are conscious but demonstrate neurologic injury such as inability to move their arms and/or legs, sensory deficits, or paresthesias.

Spinal motion restriction techniques. For these casualties, secure the individual body armor in place to protect the thoracic spine after evaluation and lifesaving interventions are performed as needed. The cervical spine may be protected by using a cervical stabilization device in conjunction with the casualty’s individual body armor or by an additional first responder holding the casualty’s head to maintain alignment with the back. Long or short spine boards should be used in addition to these measures when available.

Communication

Combat is a frightening experience. Being wounded, especially seriously, can generate tremendous anxiety and fear. Engaging a casualty with reassurance is therapeutically beneficial. Communication is just as important in casualty care on the battlefield as it is in the treatment facility. Ensure the care plan is explained to the casualty.

Documentation

Battlefield documentation of injuries and care rendered in the pre-hospital arena is sorely lacking. There is a tremendous need for documenting clinical assessments, treatment rendered, and changes in the casualty’s status, and forwarding that information with the casualty to the next level of care. Use Department of the Army (DA) Form 7656, *Tactical Combat Casualty Care Card*, for this purpose (see Figure 1-4). If this form is not available, use 3-inch white tape on the casualty’s chest and an indelible pen to document care.

The form is titled "Front" and "Back". It contains the following fields and sections:

- Name/Unit: _____
- ALLERGIES: _____
- Priority: _____ Unknown: _____ NBC: _____
- TO/TIME: _____
- Two diagrams of a human figure for marking injuries.
- A: Insect Adjunct Crit; Intubated: _____
- B: Chest Seal NeedleD Clov/Tub: _____
- C: TX Hemostatic Packed Pressure/Dress: _____
- FLUIDS: IV IO: _____
- NS/LR 500 1000 1500: _____
- Wet/Dry 500 1000: _____
- Other: _____
- DRUGS (Type / Dose / Route): _____
- PAIN: _____
- AIR: _____
- OTHER: _____
- GSW BLAST MVA Other: _____
- TIME: _____
- AVPU: _____
- PULSE: _____
- RESP: _____
- SB: _____
- First Responder's Name: _____

Figure 1-4. DA Form 7656, *Tactical Combat Casualty Care Card*

Section IV: Tactical Evacuation Care

Tactical evacuation care is the care rendered once the casualty has been picked up by an aircraft, vehicle, or boat for transportation to a higher echelon of care. Tactical evacuation is the evacuation of combat casualties from the battlefield. In general, tactical evacuation care is a continuation of care rendered during the tactical field care phase, with minor additions based on the following conditions:

- Additional medical personnel may accompany the evacuation asset.
- Additional medical equipment may be brought with the evacuation asset.

The arrival of additional medical personnel is important for several reasons:

- The medic may be one of the casualties or be dehydrated, hypothermic, or otherwise debilitated.
- There may be multiple casualties that exceed the medic's capability to care for simultaneously.
- The evacuation asset's equipment will need to be prepared prior to evacuation.
- Additional medical personnel such as physicians and other specialists provide greater expertise.

The additional medical equipment brought by the evacuation asset serves several purposes. Medical resupply may be accomplished during this phase of care. More advanced medical equipment such as blood products and other fluids, electronic monitoring devices, and oxygen may now be used to care for the casualty. This equipment and the possibly improved care environment of the evacuation asset allow more advanced casualty care with more skilled providers during the tactical evacuation phase.

Airway Management

Airway management during the tactical evacuation phase follows the same principles as during the tactical field care phase, with the use of positioning and an NPA as the initial management options. However, the management of an impaired airway is exceedingly difficult during tactical evacuation. It is now appropriate, if the equipment and provider expertise are available, to obtain a more definitive airway if required by the casualty's condition. Possible airway management options include:

Cricothyroidotomy. As in the tactical field care phase, cricothyroidotomy is still an appropriate option when an NPA is not effective. This is still the

procedure of choice for penetrating wounds of the face or neck, where blood or disrupted anatomy precludes good visualization of the vocal cords.

Intubation. The conditions of the tactical evacuation phase now make intubation a viable option. If the equipment is available and the care provider has the appropriate expertise, several intubation methods are possible. Blind rescue devices such as the laryngeal mask airway, intubating laryngeal mask airway (ILMA), King LT, or the combitube are recommended options. These devices provide adequate ventilation without the need for illuminated laryngoscopy; have been used effectively in the pre-hospital setting; and, in the case of the ILMA and combitube, protect the airway from aspiration. Additionally, if personnel have adequate training, endotracheal intubation is now an option.

Breathing

During the tactical evacuation phase, management of the casualty's breathing is a continuation of the interventions made during the tactical field care phase. Continue to treat penetrating chest wounds with occlusive dressing and monitor for the development of a tension PTX. Treat PTXs with needle decompression. At this phase of care, though, it may now be possible to consider additional interventions.

Chest tubes. For casualties with a tension PTX who show no improvement with needle decompression, the provider should consider inserting a chest tube. A chest tube should also be considered for casualties with PTX when a long evacuation time is anticipated, even if the initial needle decompression was successful.

Oxygen. Oxygen may be brought by the evacuation asset and now be available. Most combat casualties do not require oxygen, but it should be used in seriously injured casualties, especially in the followings circumstances:

- Low oxygen saturation by pulse oximetry.
- Injuries associated with impaired oxygenation.
- Unconscious casualties.
- Casualties with TBIs.
- Casualties in shock.
- Casualties at altitude.

Fluid Resuscitation

Several improvements in fluid resuscitation are possible in the tactical evacuation phase. Monitoring equipment brought by the evacuation asset may yield a better understanding of a casualty's fluid status and can direct resuscitation efforts. Continue resuscitation in casualties with a TBI to maintain a systolic blood pressure of at least 90 millimeters (mm) mercury (Hg). If indicated and available, packed red blood cells should be given to casualties suffering from blood loss. These blood cells will restore oxygen-carrying capacity. Resuscitation for casualties in shock can be continued with the use of Hextend (but not to exceed 1,000 ml per casualty) or LRS. Finally, LRS can now be used to treat dehydration.

Hypothermia Prevention

Hypothermia prevention becomes paramount during the tactical evacuation phase, especially if the casualty is evacuated in a helicopter. Continue to follow the hypothermia prevention principles of the tactical field care phase: minimize the casualty's exposure to the elements, replace wet clothing, and use equipment such as the HPMK. If the doors of the evacuation asset must be kept open, protect the casualty from the wind. Also, if portable fluid-warming devices such as the Thermal Angel are available, they should be used on all IV fluid sites.

Monitoring

The evacuation asset may contain additional patient-monitoring devices. Electronic systems capable of monitoring blood pressure, heart rate, pulse oximetry, and end-tidal carbon dioxide may be available and should be used. This is especially true in helicopter evacuation, which impairs or prevents the ability to monitor the casualty clinically.

Pneumatic Anti-Shock Garment

The pneumatic anti-shock garment (PASG) may be available during the CASEVAC phase and may be useful for stabilizing pelvic fractures and helping to control pelvic and abdominal bleeding. However, its application and extended use must be carefully monitored. This device is contraindicated and should not be used in casualties with thoracic and brain injuries.

Additional Measures

All other aspects of care during the CASEVAC phase are identical to those during the tactical field care phase. Hemorrhage must be controlled, using tourniquets as necessary. However, tourniquets should be discontinued, if possible, once bleeding is controlled by other means and the casualty has been resuscitated for hemorrhagic shock. Maintain vascular access with

at least one 18-gauge IV or an IO device, if necessary. Provide analgesia and antibiotics as previously indicated during the tactical field care phase. Continue to document all care, and forward this information with the casualty to the next level of care.

Section V: Management Guidelines

Basic Management Plan for Care under Fire

1. Return fire and take cover.
2. Direct or expect the casualty to remain engaged as a combatant if appropriate.
3. Direct the casualty to move to cover and apply self-aid if able.
4. Try to keep the casualty from sustaining additional wounds.
5. Casualties should be extricated from burning vehicles or buildings and moved to places of relative safety. Do what is necessary to stop the burning process.
6. Airway management is generally best deferred until the tactical field care phase.
7. Stop life-threatening external hemorrhage if tactically feasible.
 - Direct the casualty to control the hemorrhage by self-aid if able.
 - Use a Committee on Tactical Combat Care (CoTCCC)-recommended tourniquet for hemorrhage that is anatomically amenable to tourniquet application.
 - Apply the tourniquet proximal to the bleeding site, over the uniform, tighten, and move the casualty to cover.

Basic Management Plan for Tactical Field Care

1. Casualties with an altered mental status should be disarmed immediately.
2. Airway management.
 - Unconscious casualty without airway obstruction:
 - Chin lift or jaw-thrust maneuver.
 - NPA.
 - Place casualty in the recovery position.

- Casualty with airway obstruction or impending airway obstruction:
 - Chin lift or jaw-thrust maneuver.
 - NPA.
 - Allow casualty to assume any position that best protects the airway, to include sitting up.
 - Place unconscious casualty in the recovery position.
 - If previous measures unsuccessful, perform surgical cricothyroidotomy (with lidocaine if conscious).

3. Breathing.

- In a casualty with progressive respiratory distress and known or suspected torso trauma, consider a tension pneumothorax and decompress the chest on the side of the injury with a 14-gauge, 3.25-inch needle/catheter unit inserted in the second intercostal space at the midclavicular line. Ensure the needle entry into the chest is not medial to the nipple line and is not directed toward the heart.
- All open and/or sucking chest wounds should be treated by immediately applying an occlusive material to cover the defect and securing it in place. Monitor the casualty for the potential development of a subsequent tension pneumothorax.

4. Bleeding.

- Assess for unrecognized hemorrhage and control all sources of bleeding. If not already done, use a CoTCCC-recommended tourniquet to control life-threatening external hemorrhage that is anatomically amenable to tourniquet application or for any traumatic amputation. Apply directly to the skin 2 to 3 inches above the wound.
- For compressible hemorrhage not amenable to tourniquet use or as an adjunct to tourniquet removal (if evacuation time is anticipated to be longer than two hours), use combat gauze as the hemostatic agent of choice. Combat gauze should be applied with at least three minutes of direct pressure. Before releasing any tourniquet on a casualty who has been resuscitated for hemorrhagic shock, ensure a positive response to resuscitation efforts (i.e., a peripheral pulse normal in character and normal mentation if there is no TBI). If a lower extremity wound is not amenable to tourniquet application and cannot be controlled by hemostatics/dressings, consider immediate application of mechanical direct pressure including CoTCCC-recommended devices such as the combat ready clamp.

- Reassess prior tourniquet application. Expose wound and determine if tourniquet is needed. If so, move tourniquet from over uniform and apply directly to skin 2 to 3 inches above the wound. If a tourniquet is not needed, use other techniques to control the bleeding.
- When time and the tactical situation permit, a distal pulse check should be accomplished. If a distal pulse is still present, consider additional tightening of the tourniquet or the use of a second tourniquet, side by side and proximal to the first, to eliminate the distal pulse.
- Expose and clearly mark all tourniquet sites with the time of tourniquet application. Use an indelible marker.

5. IV access.

- Start an 18-gauge IV or saline lock if indicated.
- If resuscitation is required and IV access is not obtainable, use the IO route.

6. Tranexamic acid (TXA). If a casualty is anticipated to need significant blood transfusion (for example, presents with hemorrhagic shock, one or more major amputations, penetrating torso trauma, or evidence of severe bleeding):

- Administer 1 g of tranexamic acid in 100 cc normal saline or LRS as soon as possible but NOT later than 3 hours after injury.
- Begin second infusion of 1 g TXA after Hextend or other fluid treatment.

7. Fluid resuscitation. Assess for hemorrhagic shock. Altered mental status (in the absence of head injury) and weak or absent peripheral pulses are the best field indicators of shock.

- If not in shock:
 - No IV fluids necessary.
 - PO fluids permissible if conscious and can swallow.
- If in shock:
 - Hextend, 500-ml IV bolus.
 - Repeat once after 30 minutes if still in shock.
 - No more than 1,000 ml of Hextend.

Continued efforts to resuscitate must be weighed against logistical and tactical considerations and the risk of incurring further casualties.

If a casualty with an altered mental status due to a suspected TBI has a weak or absent peripheral pulse, resuscitate as necessary to maintain a palpable radial pulse.

8. Prevention of hypothermia.

- Minimize the casualty's exposure to the elements. Keep protective gear on or with the casualty if feasible.
- Replace wet clothing with dry clothing if possible. Get the casualty onto an insulated surface as soon as possible.
- Apply the Ready-Heat blanket from the HPMK to the casualty's torso (not directly on the skin) and cover the casualty with the heat-reflective shell.
- If a heat-reflective shell is not available, the previously recommended combination of the blizzard survival blanket and the Ready-Heat blanket may also be used.
- If the items mentioned above are not available, use dry blankets, poncho liners, sleeping bags, or anything that will retain heat and keep the casualty dry.
- Warm fluids are preferred if IV fluids are required.

9. Penetrating eye trauma. If a penetrating eye injury is noted or suspected:

- Perform a rapid field test of visual acuity.
- Cover the eye with a rigid eye shield (NOT a pressure patch). Ensure that the 400-mg Moxifloxacin tablet in the combat pill pack is taken if possible and that IV/IM antibiotics are given as outlined below if oral Moxifloxacin cannot be taken.

10. Monitoring.

- Pulse oximetry should be available as an adjunct to clinical monitoring.
- Readings may be misleading in the settings of shock or marked hypothermia.

11. Inspect and dress known wounds.

12. Check for additional wounds.

13. Provide analgesia as necessary.

- Able to fight: (**Note:** These medications should be carried by the combatant and self-administered as soon as possible after the wound is sustained.)
 - Mobic, 15 mg PO once a day.
 - Tylenol, 650-mg bilayer caplet, two PO every 8 hours.
 - Unable to fight: (**Note:** Have Naloxone readily available whenever administering opiates.)
 - Does not otherwise require IV/IO access
 - Oral transmucosal fentanyl citrate, 800 mg transbuccally:
 - * Recommend taping lozenge-on-a-stick to casualty's finger as an added safety measure.
 - * Reassess in 15 minutes.
 - * Add second lozenge, in other cheek, as necessary to control severe pain.
 - * Monitor for respiratory depression.
 - IV or IO access obtained:
 - Morphine sulfate, 5 mg IV/IO:
 - * Reassess in 10 minutes.
 - * Repeat dose every 10 minutes as necessary to control severe pain.
 - * Monitor for respiratory depression.
 - Promethazine, 25 mg IV/IM/IO:
 - * Every 6 hours as needed for nausea or for synergistic analgesic effect.
14. Splint fractures and recheck pulse.
15. Antibiotics: Recommended for all open combat wounds.
- If able to take PO:
 - Moxifloxacin, 400 mg PO, once a day.

- If unable to take PO (shock, unconsciousness):
 - Cefotetan, 2 g IV (slow push over 3 to 5 minutes) or IM every 12 hours, or
 - Ertapenem, 1 g IV/IM once a day.

16. Burns.

- Facial burns, especially those that occur in closed spaces, may be associated with inhalation injury. Aggressively monitor airway status and oxygen saturation in such patients and consider early surgical airway for respiratory distress or oxygen desaturation.
- Estimate TBSA burned to the nearest 10 percent using the “Rule of Nine.”
- Cover the burn area with dry, sterile dressings. For extensive burns (greater than 20 percent), consider placing the casualty in the HPMK to both cover the burned areas and prevent hypothermia in accordance with Section III.
- Fluid resuscitation (U.S. Army Institute of Surgical Research [USAISR] Rule of Ten):
 - If burns are greater than 20 percent of TBSA, fluid resuscitation should be initiated as soon as IV/IO access is established. Resuscitation should be initiated with LRS, normal saline, or Hextend. If Hextend is used, no more than 1,000 ml should be given, followed by LRS or normal saline as needed.
 - Initial IV/IO fluid rate is calculated as %TBSA x 10cc/hour for adults weighing 40-80 kg.
 - For every 10 kg above 80 kg, increase initial rate by 100 ml/hour.
 - If hemorrhagic shock is also present, resuscitation for hemorrhagic shock takes precedence over resuscitation for burn shock. Administer IV/IO fluids per the TCCC management guidelines.
 - Analgesia in accordance with Section III may be administered to treat burn pain.
 - Pre-hospital antibiotic therapy is not indicated solely for burns, but antibiotics should be given per the TCCC management guidelines if indicated to prevent infection in penetrating wounds.
 - All TCCC interventions can be performed on or through burned skin in a burn casualty.

17. Communicate with the casualty if possible.

- Encourage; reassure.
- Explain care.

18. CPR. Resuscitation on the battlefield for victims of blast or penetrating trauma who have no pulse, no ventilations, and no other signs of life will not be successful and should not be attempted. However, casualties with torso trauma or polytrauma who have no pulse or respirations during tactical field care should have bilateral needle decompression performed to ensure they do not have a tension pneumothorax prior to discontinuation of care. The procedure is the same as described in Section III above.

19. Documentation of care. Document clinical assessments, treatments rendered, and changes in the casualty's status on DA Form 7656. Forward this information with the casualty to the next level of care.

Basic Management Plan for Tactical Evacuation Care

The term "tactical evacuation" includes both CASEVAC and medical evacuation as defined in Joint Publication 4-02, *Doctrine for Health Services Support in Joint Operations*.

1. Airway management.

- Unconscious casualty without airway obstruction:
 - Chin lift or jaw-thrust maneuver.
 - NPA.
 - Place casualty in the recovery position.
- Casualty with airway obstruction or impending airway obstruction:
 - Chin lift or jaw-thrust maneuver.
 - NPA.
 - Allow casualty to assume any position that best protects the airway, to include sitting up.
 - Place unconscious casualty in the recovery position.
 - If above measures unsuccessful:
 - * Laryngeal mask airway (LMA)/intubating LMA.
 - * Combitube.

- * Endotracheal intubation.
- * Surgical cricothyroidotomy (with lidocaine if conscious).

- Spinal immobilization is not necessary for casualties with penetrating trauma.

2. Breathing.

- In a casualty with progressive respiratory distress and known or suspected torso trauma, consider a tension pneumothorax and decompress the chest on the side of the injury with a 14-gauge, 3.25-inch needle/catheter unit inserted in the second intercostal space at the mid-clavicular line. Ensure that the needle entry into the chest is not medial to the nipple line and is not directed toward the heart.
- Consider chest tube insertion if no improvement and/or long transport is anticipated.
- Most combat casualties do not require supplemental oxygen, but administration of oxygen may be of benefit for the following types of casualties:
 - Low oxygen saturation by pulse oximetry.
 - Injuries associated with impaired oxygenation.
 - Unconscious casualty.
 - Casualty with TBI (maintain oxygen saturation greater than 90 percent).
 - Casualty in shock.
 - Casualty at altitude.
- All open and/or sucking chest wounds should be treated by immediately applying an occlusive material to cover the defect and securing it in place. Monitor the casualty for the potential development of a subsequent tension pneumothorax.

3. Bleeding.

- Assess for unrecognized hemorrhage and control all sources of bleeding. If not already done, use a CoTCCC-recommended tourniquet to control life-threatening external hemorrhage that is anatomically amenable to tourniquet application or for any traumatic amputation. Apply directly to the skin 2 to 3 inches above the wound.

- For compressible hemorrhage not amenable to tourniquet use or as an adjunct to tourniquet removal (if evacuation time is anticipated to be longer than two hours), use combat gauze as the hemostatic agent of choice. Combat gauze should be applied with at least three minutes of direct pressure. Before releasing any tourniquet on a casualty who has been resuscitated for hemorrhagic shock, ensure a positive response to resuscitation efforts (i.e., a peripheral pulse normal in character and normal mentation if there is no TBI.) If a lower extremity wound is not amenable to tourniquet application and cannot be controlled by hemostatics/dressings, consider immediate application of mechanical direct pressure, including CoTCCC-recommended devices such as the combat ready clamp.
 - Reassess prior tourniquet application. Expose the wound and determine if a tourniquet is needed. If so, move the tourniquet from over the uniform and apply directly to the skin 2 to 3 inches above the wound. If a tourniquet is not needed, use other techniques to control the bleeding.
 - When time and the tactical situation permit, a distal pulse check should be accomplished. If a distal pulse is still present, consider additional tightening of the tourniquet or the use of a second tourniquet, side by side and proximal to the first, to eliminate the distal pulse.
 - Expose and clearly mark all tourniquet sites with the time of tourniquet application. Use an indelible marker.
4. IV access. Reassess need for IV access:
- If indicated, start an 18-gauge IV or saline lock.
 - If resuscitation is required and IV access is not obtainable, use IO route.
5. TXA. If a casualty is anticipated to need a significant blood transfusion (for example, presents with hemorrhagic shock, one or more major amputations, penetrating torso trauma, or evidence of severe bleeding):
- Administer 1 g of tranexamic acid in 100 cc normal saline or LRS as soon as possible but NOT later than 3 hours after injury.
 - Begin second infusion of 1 g TXA after Hextend or other fluid treatment.
6. Fluid resuscitation. Reassess for hemorrhagic shock (altered mental status in the absence of brain injury and/or change in pulse character). If blood pressure monitoring is available, maintain target systolic blood pressure 80-90 mmHg.

- If not in shock:
 - No IV fluids necessary.
 - PO fluids permissible if conscious and can swallow.
- If in shock and blood products are not available:
 - Hextend 500-ml IV bolus.
 - Repeat after 30 minutes if still in shock.
 - Continue resuscitation with Hextend or crystalloid solution as needed to maintain target blood pressure or clinical improvement.
- If in shock and blood products are available under an approved command or theater protocol:
 - Resuscitate with two units of plasma followed by packed red blood cells in a 1:1 ratio. If blood component therapy is not available, transfuse fresh whole blood. Continue resuscitation as needed to maintain target blood pressure or clinical improvement.
- If a casualty with an altered mental status due to a suspected TBI has a weak or absent peripheral pulse, resuscitate as necessary to maintain a palpable radial pulse. If blood pressure monitoring is available, maintain target systolic blood pressure of at least 90 mmHg.

7. Prevention of hypothermia.

- Minimize the casualty's exposure to the elements. Keep protective gear on or with the casualty if feasible.
- Replace wet clothing with dry if possible. Get the casualty onto an insulated surface as soon as possible.
- Apply the Ready-Heat blanket from the HPMK to the casualty's torso (not directly on the skin), and cover the casualty with the heat-reflective shell.
- If a heat-reflective shell is not available, the previously recommended combination of the blizzard survival blanket and the Ready-Heat blanket may also be used.
- If the items mentioned above are not available, use poncho liners, sleeping bags, or anything that will retain heat and keep the casualty dry.
- Use a portable fluid warmer capable of warming all IV fluids, including blood products.

- Protect the casualty from the wind if the doors to the evacuation asset must be kept open.
8. Penetrating eye trauma. If a penetrating eye injury is noted or suspected:
- Perform a rapid field test of visual acuity.
 - Cover the eye with a rigid eye shield (NOT a pressure patch).
 - Ensure the 400-mg Moxifloxacin tablet in the combat pill pack is taken if possible and that IV/IM antibiotics are given as outlined below if oral Moxifloxacin cannot be taken.
9. Monitoring. Institute pulse oximetry and other electronic monitoring of vital signs, if indicated.
10. Inspect and dress known wounds if not already done.
11. Check for additional wounds.
12. Provide analgesia as necessary.
- Able to fight: (**Note:** These medications should be carried by the combatant and self-administered as soon as possible after the wound is sustained.)
 - Mobic, 15 mg PO once a day.
 - Tylenol, 650-mg bilayer caplet, two PO every 8 hours.
 - Unable to fight: (**Note:** Have Naloxone readily available whenever administering opiates.)
 - Does not otherwise require IV/IO access:
 - Oral transmucosal fentanyl citrate (OTFC), 800 mg transbuccally:
 - * Recommend taping lozenge-on-a-stick to casualty's finger as an added safety measure.
 - * Reassess in 15 minutes.
 - * Add second lozenge, in other cheek, as necessary to control severe pain.
 - * Monitor for respiratory depression.
 - IV or IO access obtained:
 - Morphine sulfate, 5 mg IV/IO.
 - * Reassess in 10 minutes.

- * Repeat dose every 10 minutes as necessary to control severe pain.
 - * Monitor for respiratory depression.
 - Promethazine, 25 mg IV/IM/IO:
 - * Every 6 hours as needed for nausea or for synergistic analgesic effect.
13. Reassess fractures and recheck pulses.
14. Antibiotics: Recommended for all open combat wounds.
- If unable to take PO (shock, unconsciousness):
 - Cefotetan, 2 g IV (slow push over 3 to 5 minutes) or IM every 12 hours, or
 - Ertapenem, 1 g IV/IM once a day.
15. Burns.
- Facial burns, especially those that occur in closed spaces, may be associated with inhalation injury. Aggressively monitor airway status and oxygen saturation in such patients and consider early surgical airway for respiratory distress or oxygen desaturation.
 - Estimate TBSA burned to the nearest 10 percent using the Rule of Nine.
 - Cover the burn area with dry, sterile dressings. For extensive burns (greater than 20 percent), consider placing the casualty in the HPMK to both cover the burned areas and prevent hypothermia in accordance with Section III.
 - Fluid resuscitation (USAISR Rule of Ten):
 - If burns are greater than 20 percent of the TBSA, fluid resuscitation should be initiated as soon as IV/IO access is established. Resuscitation should be initiated with LRS, normal saline, or Hextend. If Hextend is used, no more than 1,000 ml should be given, followed by LRS or normal saline as needed.
 - Initial IV/IO fluid rate is calculated as %TBSA x 10cc/hour for adults weighing 40-80 kg.
 - For every 10 kg above 80 kg, increase initial rate by 100 ml/hour.

- If hemorrhagic shock is also present, resuscitation for hemorrhagic shock takes precedence over resuscitation for burn shock. Administer IV/IO fluids per the TCCC guidelines.

- Analgesia in accordance with Section III may be administered to treat burn pain.
- Pre-hospital antibiotic therapy is not indicated solely for burns, but antibiotics should be given per the TCCC guidelines if indicated to prevent infection in penetrating wounds.
- All TCCC interventions can be performed on or through burned skin in a burn casualty.

16. The PASG may be useful for stabilizing pelvic fractures and controlling pelvic and abdominal bleeding. Application and extended use must be carefully monitored. The PASG is contraindicated for casualties with thoracic or brain injuries.

17. CPR in tactical evacuation care.

- Casualties with torso trauma or polytrauma who have no pulse or respirations during tactical evacuation should have bilateral needle decompression performed to ensure they do not have a tension pneumothorax. The procedure is the same as described in Section III above.
- CPR may be attempted during this phase of care if the casualty does not have obviously fatal wounds and will be arriving at a facility with a surgical capability within a short period of time. CPR should not be done at the expense of compromising the mission or denying lifesaving care to other casualties.

18. Documentation of care. Document clinical assessments, treatments rendered, and changes in casualty's status on DA Form 7656. Forward this information with the casualty to the next level of care.

Chapter 2

Tactical Combat Casualty Care Procedures

Section I: Hemorrhage Control

Combat Application Tourniquet

(Necessary equipment: Combat application tourniquet [CAT])



Figure 2-1. CAT



Figure 2-2

1. Place the wounded extremity through the loop of the band. Position the tourniquet about 2 inches above the injury site.



Figure 2-3

2. Route the self-adhering band through both sides of the friction adapter buckle.



Figure 2-4

3. Feed the self-adhering band tight around the extremity and securely fasten it back on itself.



Figure 2-5

4. Adhere it completely around the band until it reaches the clip.



Figure 2-6

5. Twist the windlass rod until the bleeding stops.



Figure 2-7

6. Lock the windlass rod in place with the windlass clip.



Figure 2-8

7. Grasp the windlass strap, pull tight, and adhere it to the windlass clip.

8. If the tactical situation permits, check for a distal pulse. If a distal pulse is still present, apply a second tourniquet side by side and proximal to the first. Tighten this tourniquet and recheck the distal pulse.

Note: During self-application of the CAT to an upper extremity wound, defer routing the self-adhering band through both sides of the friction-adaptor buckle.

Special Operations Forces Tactical Tourniquet

(Necessary equipment: Special operations forces tactical tourniquet [SOFTT])



Figure 2-9. SOFTT



Figure 2-10

1. Place the wounded extremity through the loop of the band, positioning the tourniquet about 2 inches above the injury site.



Figure 2-11

2. Pull the tail tight ensuring to remove as much slack as possible.



Figure 2-12

3. Twist set screw clockwise to lock clamp in place.



Figure 2-13

4. Twist the windlass rod until the bleeding stops.



Figure 2-14

5. Lock the windlass rod in place with the windlass tri-ring anchor. It is not necessary to lock both ends of the windlass.

Combat Gauze

(Necessary equipment: Combat gauze)

1. Apply dressing with pressure to the wound for 3 minutes.
2. If bleeding continues after 3 minutes of pressure, remove first combat gauze and repeat step 1.
3. Once bleeding is controlled, apply outer bandage (Ace wrap or emergency dressing) to secure the dressing to the wound.

Section II: Airway Management

Nasopharyngeal Airway Insertion

(Necessary equipment: Nasopharyngeal airway [NPA], gloves, water-based lubricant)

1. Place the casualty supine with the head in a neutral position.

Caution: Do not use the NPA if there is clear fluid (cerebrospinal fluid) coming from the ears or nose. This may indicate a skull fracture.

2. Select the appropriately sized airway using one of the following methods:

- Measure the airway from the casualty's nostril to the earlobe.
- Measure the airway from the casualty's nostril to the angle of the jaw.

Note: Choosing the proper length ensures appropriate diameter. Standard adult sizes are 34, 32, 30, and 28 French (Fr).

3. Lubricate the tube with a water-based lubricant.

Caution: Do not use a petroleum-based or non-water-based lubricant. These substances can cause damage to the tissues lining the nasal cavity and pharynx, increasing the risk for infection.

4. Insert the NPA.

- Push the tip of the nose upward gently.
- Position the tube so the bevel of the airway faces toward the septum.
- Insert the airway into the nostril and advance it until the flange rests against the nostril.

Caution: Never force the NPA into the casualty's nostril. If resistance is met, pull the tube out and attempt to insert it in the other nostril. Most attempts to insert the NPA should be in the right nostril. If unable to insert into the right nostril, try the left. If inserting in the left nostril, the bevel will not be against the septum.




Figure 2-15. NPA insertion

Surgical Cricothyroidotomy

(Necessary equipment: Prefabricated cricothyroidotomy kit. In the event kits are unavailable, an improvised kit should include a cutting instrument [for example, Scalpel #10 or #15]; forceps or tracheal hook; povidone-iodine; endotracheal tube [ETT], 6 millimeter [mm]; gloves; 4 x 4 gauze; tape; local anesthetic; and materials to inject)

Note: Cricothyroidotomy sets should be prepared prior to the mission. All essential pieces of equipment should be prepared before deployment and packed into a Ziploc bag. Cut the ETT to just above the cuff inflation tube so the ETT is not protruding 6 inches out of the casualty's neck.




Figure 2-16. Surgical airway (cricothyroidotomy)

1. Hyperextend the casualty's neck.
 - Place the casualty in the supine position.
 - Place a blanket or poncho rolled up under the casualty's neck or between the shoulder blades so the airway is straight.

Warning: Do not hyperextend the casualty's neck if a cervical injury is suspected.

2. Put on medical gloves, available in patient's individual first aid kit.
3. Locate the cricothyroid membrane.
 - Place a finger of the nondominant hand on the thyroid cartilage (Adam's apple), and slide the finger down to the cricoid cartilage.
 - Palpate for the "V" notch of the thyroid cartilage.
 - Slide the index finger down into the depression between the thyroid and cricoid cartilage.




Figure 2-17. Cricothyroid membrane anatomy

4. Prepare the incision site.
 - Administer local anesthesia to the incision site if the casualty is conscious.
 - Prep the skin over the membrane with an alcohol pad or povidone-iodine.
5. With a cutting instrument in the dominant hand, make a 1.5-inch vertical incision through the skin over the cricothyroid membrane.

Caution: Do not cut the cricothyroid membrane with this incision.

6. Relocate the cricothyroid membrane by touch and sight.
7. Stabilize the larynx with one hand, and make a 1/2-inch horizontal incision through the elastic tissue of the cricothyroid membrane.

Note: A rush of air may be felt through the opening.

8. Dilate the opening with a hemostat or scalpel handle. Hook the cricothyroid membrane with a prefabricated cricothyroid hook or bent 18-gauge needle.

9. Grasp the cricoid cartilage and stabilize it.

10. Insert the ETT through the opening and toward the lungs. Only advance the ETT 2 to 3 inches into the trachea to prevent right main stem bronchus intubation. Inflate the cuff to prevent aspiration.

11. Secure the tube circumferentially around the patient's neck to prevent accidental extubation. This can be achieved with tape, tubing, or a prefabricated device in some kits.

12. Check for air exchange and tube placement.

- Air exchange: Listen and feel for air passage through the tube; look for fogging in the tube.
- Tube placement: Bilateral chest sounds/rise and fall of the chest confirm proper tube placement.
- Unilateral breath sounds/rise and fall of chest indicate a right main stem bronchus intubation. Withdraw the ETT 1 to 2 inches and reconfirm placement.
- Air from the casualty's mouth indicates the tube is directed toward the mouth. Remove the tube, reinsert, and recheck for air exchange and placement.
- Any other problem indicates the tube is not placed correctly. Remove the tube, reinsert, and recheck for air exchange.

13. Once the tube is correctly placed, begin rescue breathing, if necessary and tactically appropriate.

- Connect the tube to a bag valve mask and ventilate the casualty at the rate of 20 breaths/minute.
- If a bag valve mask is not available, begin mouth-to-tube resuscitation at 20 breaths/minute.

14. Apply a sterile dressing. Use either of the following methods:

- Make a V-shaped fold in a 4 x 4 gauze pad and place it under the edge of the ETT to prevent irritation to the casualty. Tape securely.
- Cut two 4 x 4 gauze pads halfway through and place on opposite sides of the tube. Tape securely.

King LT Insertion

(Necessary equipment: King LT, water-based lubricant, syringe)

1. Prepare the casualty.
 - Place the casualty’s head in the “sniffing” position.
 - Preoxygenate the casualty, if equipment is available.
2. Prepare the King LT.
 - Choose the appropriately sized tube.
 - Test cuff inflation by injecting the proper volume of air into the cuff. Deflate the cuff prior to inserting the tube.
 - Lubricate the tube with a water-based lubricant.

Caution: Do not use a petroleum-based or non-water-based lubricant. These substances can cause damage to the tissues lining the nasal cavity and pharynx, increasing the risk for infection.

3. Insert the King LT.
 - Hold the tube in the dominant hand. With the nondominant hand, open the casualty’s mouth and apply a chin lift.
 - With the King LT rotated laterally 45 to 90 degrees, place the tip into the mouth and advance the tube behind the base of the tongue.

Note: A lateral approach with the chin lift facilitates proper insertion. The tip must remain midline as it enters the posterior pharynx.

- Rotate the tube to midline as the tip reaches the posterior pharynx.
 - Advance the tube until the base of the connector is aligned with the teeth or gums.
 - Using either an attached pressure gauge or syringe, inflate the cuff to the minimum volume necessary to seal the airway.
4. Confirm proper placement of the tube.
 - Reference marks for the tube are at the proximal end of the tube and should be aligned with the upper teeth.
 - Confirm proper placement by listening for equal breath sounds during ventilation.

- While gently ventilating the casualty, withdraw the tube until ventilation is easy and free flowing, with minimal airway pressure needed.

Note: Initially placing the tube deeper than required and then withdrawing slightly increases the chance of proper insertion, helps ensure a patient airway, and decreases the risk of airway obstruction if the casualty spontaneously ventilates.

5. Secure the tube with tape.

Section III: Breathing Management

Penetrating Chest Wounds

(Necessary equipment: Field dressings or any airtight material [occlusive chest seal, plastic wrap])

1. Expose the wound(s).

- Cut or unfasten the clothing that covers the wound.
- Wipe blood/sweat from skin surrounding wound to increase seal effectiveness.
- Disrupt the wound as little as possible.

Note: Do not remove clothing stuck to the wound.

2. Check for an exit wound.

- Feel and/or look at the casualty's chest and back.
- Remove the casualty's clothing, if necessary.

3. Seal the wound(s), usually covering the first wound encountered.

Note: All penetrating chest wounds should be treated as if they are sucking chest wounds.

Note: In an emergency, any airtight material can be used. The material must be large enough so it is not sucked into the chest cavity.

- Cut the dressing wrapper on one long and two short sides and remove the dressing.
- Apply the inner surface of the wrapper to the wound when the casualty exhales.
- Ensure that the covering extends at least 2 inches beyond the edges of the wound.

- Seal by applying overlapping strips of tape to all edges of occlusive dressing, forming a complete seal.
 - Cover the exit wound in the same way, if applicable.
4. Place the casualty on the injured side or sitting up.
 5. Monitor the casualty.
 - Monitor breathing and the wound seal for continued effectiveness.
 - Check vital signs.
 - Observe for signs of shock.

Needle Chest Decompression

(Necessary equipment: Large-bore needle [10- to 14-gauge], at least 3.25 inches in length, and tape)

1. Locate the second intercostal space (between the second and third ribs) at the midclavicular line (approximately in line with the nipple) on the affected side of the casualty's chest.
2. Insert a large-bore (10- to 14-gauge) needle/catheter unit.
 - Place the needle tip on the insertion site (second intercostal space, midclavicular line).
 - Lower the proximal end of the needle to permit the tip to enter the skin just above the third rib margin.
 - Firmly insert the needle into the skin over the third rib at a 90-degree angle to the chest wall until the pleura has been penetrated, as evidenced by feeling a "pop" as the needle enters the pleural space and a hiss of air escapes from the chest.

Warning: Proper positioning of the needle is essential to avoid puncturing blood vessels and/or nerves. The needle should not be inserted medial to the nipple line, as this will cause the needle to enter the cardiac box.

Note: If you are using a catheter-over-needle, the needle catheter should be inserted to the hub. Withdraw the needle along the angle of insertion while holding the catheter still.

3. Secure the catheter to the chest with tape, and monitor the casualty for possible return of symptoms.

Warning: Although leaving the catheter in place will reduce the likeliness of a tension pneumothorax, the patient may still develop symptoms if the catheter becomes kinked or plugged under the skin.




Figure 2-18. Needle chest compression, needle insertion site

Chest Tube Insertion

(Necessary equipment: Chest tube [16-35 Fr], gloves, one-way valve, scalpel handle and blades [#10 and #15], Kelly forceps, large hemostat, povidone-iodine, suture material, lidocaine with 1 percent epinephrine for injection, needle, and syringe)

1. Assess the casualty.
 - If necessary, open the airway.
 - Ensure adequate respiration and assist as necessary.
 - Provide supplemental oxygen, if available.
 - Connect the casualty to a pulse oximeter, if available.
2. Prepare the casualty.
 - Place the casualty in the supine position.
 - Raise the arm on the affected side above the casualty's head.
 - Select the insertion site at the anterior axillary line over the fourth or fifth intercostal space.
 - Clean the site with povidone-iodine solution.
 - Put on sterile gloves.
 - Drape the area.

- Liberally infiltrate the area with the 1 or 2 percent lidocaine solution and allow time for medication to take effect if patient symptoms permit.

3. Insert the tube.

- Make a 2 to 3 centimeter (cm) transverse incision over the selected site and extend it down to the intercostal muscles.

Note: The skin incision should be 1 to 2 cm below the intercostal space through which the tube will be placed.

- Insert the large forceps through the intercostal muscles in the next intercostal space above the skin incision.
- Puncture the parietal pleura with the tip of the forceps and slightly enlarge the hole by opening the clamp 1.5 to 2 cm.

Caution: Avoid puncturing the lung. Always use the superior margin of the rib to avoid the intercostal nerves and vessels.

- Immediately insert a gloved finger in the incision to clear any adhesions, clots, etc.
- Grasp the tip of the chest tube with forceps. Insert the tip of the tube into the incision as you withdraw your finger.
- Advance the tube until the last side hole is 2.5 to 5 cm inside the chest wall.
- Connect the end of the tube to a one-way drainage valve (e.g., Heimlich valve or improvised).
- Secure the tube using the suture materials.
- Apply an occlusive dressing over the incision site.
- Radiograph the chest to confirm placement, if available.

4. Reassess the casualty.

- Check for bilateral breath sounds.
- Monitor and record vital signs every 15 minutes.

5. Document the procedure.




Figure 2-19. Chest tube insertion site

Section IV: Vascular Access

Peripheral Intravascular Access

(Necessary equipment: IV tubing, IV fluids, 18-gauge or larger IV needle/catheter, constricting band, antiseptic wipes, gloves, tape, 2 x 2 gauze sponges)

1. Select an appropriate access site on an extremity.
 - Avoid sites over joints.
 - Avoid injured extremities.
 - Avoid extremities with significant wounds proximal to the IV insertion site.
2. Prepare the site.
 - Apply the constriction band around the limb, about 2 inches above the puncture site.
 - Clean the site with antiseptic solution.
3. Put on gloves.
4. Puncture the vein.
 - Stabilize skin at the puncture site with nondominant thumb, pulling the skin downward until taut. Avoid placing thumb directly over the vein to avoid collapsing.
 - Position the needle point, bevel up, parallel to the vein, 1/2 inch below the venipuncture site.
 - Hold the needle at a 20- to 30-degree angle and insert it through the skin.

- Move the needle forward about 1/2 inch into the vein.
- Confirm the puncture by observing blood in the flash chamber.

Note: A faint give may be felt as the needle enters the vein.

5. Advance the catheter.

- Grasp the hub and advance the needle into the vein up to the hub.

Note: This prevents backflow of blood from the hub.

- While holding the hub, press lightly on the skin with the fingers of the other hand.
- Remove the needle from the catheter and secure it in a safe place to avoid accidental needle sticks.

6. Connect the catheter to the IV infusion tubing and begin the infusion.

- If the casualty does not require IV fluids, attach a saline lock.
- Observe the site for infiltration of fluids into the tissues.

7. Secure the catheter and tubing to the skin and dress the site.

Intraosseous Placement: First Access for Shock and Trauma (FAST1) System

(Necessary equipment: First Access for Shock and Trauma (FAST1) System device, infusion fluids)

1. Prepare the site.

- Expose the sternum.
- Identify the sternal notch.

2. Place the target patch.

- Remove the top half of the backing (“Remove 1”) from the patch.
- Place index finger on the sternal notch, perpendicular to the skin.
- Align the locating notch in the target patch with the sternal notch.
- Verify that the target zone (circular hole) of the patch is directly over the casualty’s midline, and press firmly on the patch to engage the adhesive and secure the patch.
- Remove the remaining backing (“Remove 2”) and secure the patch to the casualty.



Figure 2-20. FAST1 target patch

3. Insert the introducer.

- Position yourself over the head of the patient facing toward the patient's feet.
- Remove the cap from the introducer.
- Place the bone probe cluster needles in the target zone of the target patch.
- Hold the introducer perpendicular to the skin of the casualty.
- Pressing straight along the introducer axis, with hand and elbow in line, push with a firm, constant force until a release is heard and felt.
- Expose the infusion tube by gently withdrawing the introducer. The stylet supports will fall away.
- Locate the orange sharps plug. Place it on a flat surface with the foam facing up, and keeping both hands behind the needles, push the bone probe cluster into the foam.




Figure 2-21. FAST1 introducer insertion

Warning: Avoid extreme force or twisting and jabbing motions.

4. Connect the infusion tube.
 - Connect the infusion tube to the right-angle female connector.
 - Flush catheter with 1 milliliter (ml) of sterile IV solution.
 - Attach the straight female connector to the source of fluids or drugs.



Figure 2-22. Secure the protector dome.

5. Place the protector dome directly over the target patch and press firmly to engage the Velcro fastening.

Section V: Hypothermia Prevention

Tactical Field Care Phase

1. Stop bleeding and resuscitate appropriately. If warm fluids are available, use them. They likely will not be available.
2. Remove any wet clothes and replace them with dry clothes, if possible.
3. Use the hypothermia prevention and management kit (HPMK).
 - Place the casualty on the heat-reflective shell.
 - Place the Ready-Heat blanket on the casualty's torso and back. Once the ingredients are exposed to the air, they instantly start to heat up to a maximum temperature of 104 F (40 C) for 8 hours. Do not place the Ready-Heat blanket directly on the casualty's skin, as this may cause a burn.

- Wrap the heat-reflective shell around the casualty. If a heat-reflective shell is not available, use a blizzard blanket to wrap the casualty.

Note: If you do not have an HPMK or a survival blanket of any kind, then find dry blankets, poncho liners, space blankets, sleeping bags, body bags, or anything that will retain heat and keep the casualty dry.



Figure 2-23. HPMK

Tactical Evacuation Care Phase

1. The casualty should remain wrapped in the Ready-Heat blanket, heat-reflective shell, or blizzard blanket.
2. If these items were not available in the other phases of care, check with evacuation personnel to see if they have them or any other items that can be used to prevent heat loss.
3. Wrap the casualty in dry blankets and, during helicopter transport, try to keep the wind from open doors from blowing over or under the casualty.
4. Use a portable fluid warmer on all IV sites, if available.

Section VI: Medication Guidelines

Analgesia

Note: Pain medication should be given to any casualty in pain if there are not contraindications.

If able to continue to fight:

- Oral analgesia: Give to any casualty still able to fight.

Note: These medications should be included in the combat pill pack: Meloxicam (Mobic), 15 milligrams (mg); and Acetaminophen (Tylenol), 650 mg bi-layer caplet, x 2.

If unable to continue to fight:

- Morphine sulfate:
 - Give morphine sulfate, 5 mg IV or intraosseous (IO).
 - Reassess in 10 minutes.
 - Repeat dose every 10 minutes as necessary to control severe pain.
 - Monitor for respiratory depression.
- Promethazine, 25 mg IV/IO/IM, every 4 hours for synergistic analgesic effect and as a counter to morphine-induced nausea.
- Naloxone (Narcan):
 - Have Naloxone (Narcan) available before administering morphine.
 - Use immediately if casualty exhibits signs of respiratory depression.

Antibiotics

Note: Treat all open combat wounds with antibiotics as soon as situation permits. Broad-spectrum antibiotics (for example, fluoroquinolones or cephalosporins) should be used depending on the casualty's condition and allergies.

1. Oral antibiotics:

- **Note:** This medication should be included in the combat pill pack: Moxifloxacin, 400 mg once a day.

2. Non-oral antibiotics:

- Cefotetan, 2 grams (gm) IV/IO (given with a slow push over 3 to 5 minutes) or IM every 12 hours.
- Ertapenem, 1 gm IV or IM every 24 hours.

Appendix A

Triage Categories

Triage Category	Category Description	Examples
Immediate	This group includes those Soldiers requiring lifesaving surgery. The surgical procedures in this category should not be time-consuming and should concern only those patients with high chances of survival.	Upper airway obstruction Severe respiratory distress Life-threatening bleeding Tension pneumothorax Hemothorax Flail chest Extensive 2nd- or 3rd-degree burns Untreated poisoning (chemical agent) and severe symptoms Heat stroke Decompensated shock Rapidly deteriorating level of consciousness Any other life-threatening condition that is rapidly deteriorating
Delayed	This group includes those wounded who are badly in need of time-consuming surgery, but whose general condition permits delay in surgical treatment without unduly endangering life. Sustaining treatment will be required (e.g., stabilizing intravenous fluids, splinting, and administration of antibiotics, catheterization, gastric decompression, and relief of pain).	Compensated shock Fracture, dislocation, or injury causing circulatory compromise Severe bleeding, controlled by a tourniquet or other means Suspected compartment syndrome Penetrating head, neck, chest, back, or abdominal injuries without airway or breathing compromise or decompensated shock Uncomplicated immobilized cervical spine injuries Large, dirty, or crushed soft-tissue injuries Severe combat stress symptoms or psychosis

Triage Category	Category Description	Examples
Minimal	These casualties have relatively minor injuries and can effectively care for themselves or can be helped by non-medical personnel.	Uncomplicated closed fractures and dislocations Uncomplicated or minor lacerations (including those involving tendons, muscles, and nerves) Frostbite Strains and sprains Minor head injury (loss of consciousness of less than five minutes with normal mental status and equal pupils)
Expectant	Casualties in this category have wounds that are so extensive that even if they were the sole casualty and had the benefit of optimal medical resource application, their survival would be unlikely. The expectant casualty should not be abandoned, but should be separated from the view of other casualties. Using a minimal but competent staff, provide comfort measures for these casualties.	Traumatic cardiac arrest Massive brain injury 2nd-or 3rd-degree burns over 70 percent of the body surface area Gunshot wound to the head with a Glasgow Coma Scale of 3

Appendix B

Evacuation Categories

Evacuation Category	Army	Navy	Marines	Air Force
Urgent (to save life, limb, or eyesight)	Within 1 hour	Within 1 hour	Within 1 hour	As soon as possible
Priority (medical condition could deteriorate)	Within 4 hours	Within 4 hours	Within 4 hours	Within 24 hours
Routine (condition is not expected to deteriorate significantly while awaiting flight)	Within 24 hours	Within 24 hours	Within 24 hours	Within 72 hours

Note: The categories of evacuation precedence are urgent, priority, and routine. The evacuation time periods are flexible, mission-dependent, and vary greatly among the services based upon the different types of evacuation assets that each uses. The Army uses an “Urgent Surgical” subcategory to identify casualties who may need immediate surgical intervention. The Army also uses a “Convenience” category for personnel requiring medical evacuation for conditions that are not expected to significantly change for an extended period of time (greater than 72 hours).

Appendix C

9-Line Medical Evacuation

Line 1. Location of the pickup site.

Line 2. Radio frequency, call sign, and suffix.

Line 3. Number of patients by precedence:

- A - Urgent
- B - Urgent Surgical
- C - Priority
- D - Routine
- E - Convenience

Line 4. Special equipment required:

- A - None
- B - Hoist
- C - Extraction equipment
- D - Ventilator

Line 5. Number of patients:

- A - Litter
- B - Ambulatory

Line 6. Security at pickup site:*

- N - No enemy troops in area
- P - Possible enemy troops in area (approach with caution)
- E - Enemy troops in area (approach with caution)
- X - Enemy troops in area (armed escort required)

Line 7. Method of marking pickup site:

- A - Panels
- B - Pyrotechnic signal
- C - Smoke signal
- D - None
- E - Other

* In peacetime: Number and type of wounds, injuries, and illnesses (but also desired in wartime for planning purposes).

Line 8. Patient nationality and status:

- A - U.S. military
- B - U.S. civilian
- C - Non-U.S. military
- D - Non-U.S. civilian
- E - Enemy prisoner of war

Line 9. Nuclear, biological, and chemical (NBC) contamination:**

- N - Nuclear
- B - Biological
- C - Chemical

** In peacetime: Terrain description of pickup site (but also desired in wartime, as NBC contamination is rarely an issue).

Appendix D

Combat Pill Pack

Contents:

1. Meloxicam (Mobic), 15 milligrams (mg) x 1
2. Acetaminophen (Tylenol), 500 mg x 2
3. Moxifloxacin, 400 mg x 1

Instructions: In the event of an open combat wound, swallow all four pills with water.

Note: Soldiers should be instructed in the use of the combat pill pack and should be issued the pack prior to combat.

Warning: Do not issue the pill pack to those Soldiers with known drug allergies to any of the components. In these cases it will be necessary to replace the contents with appropriate substitutes.



Figure D-1. Combat pill pack

Appendix E

Improved First-Aid Kit

The improved first-aid kit is a rapid-fielding initiative item. It is issued to deploying units by the unit's central issue facility.

Contents:

- Nasopharyngeal airway.
- Exam gloves (4).
- 2-inch tape.
- Trauma dressing.
- Kerlix (gauze bandage rolls).
- Combat application tourniquet.
- Modular lightweight load-carrying equipment pouch with retaining lanyard.



Figure E-1

Appendix F

Warrior Aid and Litter Kit

The following items are included as components of the warrior aid and litter kit (WALK):

- 1 x bag (WALK)
- 10 x gloves (trauma, nitrile, Black Talon [5 pair])
- 2 x nasopharyngeal airway (28 French with lubricant)
- 2 x gauze (Petrolatum 3" x 18")
- 2 x needle/catheter (14 gauge x 3.25")
- 2 x combat application tourniquet
- 6 x trauma dressing
- 4 x gauze (compressed, vacuum-sealed)
- 1 x emergency trauma abdominal dressing
- 2 x SAM II splint
- 1 x shears (trauma, 7.25")
- 2 x tape (surgical, adhesive 2")
- 1 x card (reference, combat casualty)
- 2 x card (individual, combat casualty)
- 1 x panel (recognition, orange)
- 1 x litter (evacuation platform, Talon 90C)
- 1 x hypothermia management and prevention kit
- 4 x strap (tie down, universal litter)

Note: There should be a WALK on at least one vehicle per convoy.



Figure F-1. WALK

Appendix G

Aid Bag Considerations

There is not a standard packing list for an aid bag. The contents of a tactical provider's aid bag are dependent upon:

- The skill level of the tactical provider.
- The type of mission.
- The length of mission.

The overall approach is not the packing of a single aid bag, but developing a tiered approach and supplying it with appropriate levels of medical supplies to meet the challenges in the different stages of care. Tactical combat casualty care, and the supplies that facilitate that care, start with each Soldier's improved first-aid kit and increase in application and amount to meet mission requirements and any worst-case scenario. Attempts should be made to pack the aid bags and stage them appropriately. The specific types and abundance of medical supplies in the proper location will ensure success. Planning and packing each bag is based on mission analysis, threats, and assets available considering a worst-case scenario during the different stages of care.

Appendix H

National Stock Numbers

Equipment	National Stock Number
Airway Supplies	
Nasopharyngeal airway	6515-00-300-2900
King LT:	Size 3: 6515-01-515-0146 Size 4: 6515-01-515-0151 Size 5: 6515-01-515-0161
Emergency Cricothyrotomy Kit	6515001-573-0692
Breathing Supplies	
Bolin Chest Seal	6501-01-549-0939
Hyfin Chest Seal	6515-01-532-8019
H&H Wound Seal Kit	6510-01-573-0300
HALO Chest Seal	6515-01-532-8019
Needle Decompression (14 gauge x 3.25 in)	6515-01-541-0635
Hemorrhage Supplies	
Combat Application Tourniquet	6515-01-521-7976
Special Operations Forces Tactical Tourniquet	6515-01-530-7015
Quick Clot Combat Gauze	6510-01-562-3325
H&H Compressed Bandage Gauze (Improved First-Aid Kit [IFAK])	6510-01-503-2117
NARS Rolled Gauze (Warrior Aid and Litter Kit [WALK])	6510-01-529-1213
Emergency Dressing (IFAK)	6510-01-492-2275
Vascular Access/Fluids	
FAST1 IO System	6515-01-453-0960
EZ IO Driver	6515-01-537-9615
EZ IO Driver Needle Sets	6515-01-537-9007 (Adult) 6515-01-518-8497 (Pediatric)
EZ IO Manual Needle Set (Non-Sternal)	6515-01-540-9794
Bone Injection Gun	6515-01-518-8487 (Adult) 6515-01-518-8497 (Pediatric)
Hetastarch (Hextend) 500 ml	6505-01-498-8636
Tactical IV Starter Kit	6515-01-587-5717

CENTER FOR ARMY LESSONS LEARNED

Hypothermia Prevention	
Hypothermia Prevention and Management Kit	6515-01-532-8056
Blizzard Survival Blanket	6532-01-534-6932
Ready-Heat Blanket	6532-01-525-4062
Miscellaneous Supplies	
Sked Basic Rescue System	6530-01-260-1222
WALK	6545-01-587-1199
IFAK	6545-01-532-4962
Talon II Model 90C	6530-01-504-9051
Sam Splint II	6515-01-494-1951
Combat Eye Shield	6515-01-590-2668

Appendix I

References/Resources

Books

Bond, C. (2010). *68W Advanced Field Craft: Combat Medic Skills*. American Association of Orthopedic Surgeons. Sudbury, MA: Jones and Bartlett.

Department of Defense (2004). *Emergency War Surgery. 3rd Revision*. Borden Institute.

National Association of Emergency Medical Technicians (2011). *Prehospital Trauma Life Support Manual, Military 7th edition*. St. Louis, MO: Mosby.

Articles

Butler, F. (1996). Tactical combat casualty care in special operations. *Military Medicine*. 161(3), 3–16.

Butler, F. (2003). Antibiotics in tactical combat casualty care. *Military Medicine*. 168, 911–914.

Cotton, B. (2009). Guidelines for pre-hospital fluid resuscitation in the injured patient. *Journal of Trauma Injury, Infection and Critical Care*. 67, 389–402.

Giebner, S. (2003). Tactical combat casualty care. *Journal of Special Operations Medicine*. 3(4). 47–55.

Kragh, J. (2009). Survival with emergency tourniquet use to stop bleeding in major limb trauma. *Annals of Surgery*, 249, 1–7.

Kragh, J. (2010). Use of tourniquets and their effects on limb function in the modern combat environment. *Foot Ankle Clin N Am* 15, 23–40.

Training Manuals

STP 8-68W13-SM-TG (2009). *Soldier's Manual and Trainer's Guide: MOS 68W, Health Care Specialist*.

Center for Army Lessons Learned (CALL) Resources

CALL Newsletter 04-18, *Medical Planning*.

CALL Special Edition 05-8, *Deploying Health Care Provider*.

Online Resources

U.S. Army Medical Department (AMEDD) Center and School Portal,
Deployment Relevant Training: [https://www.cs.amedd.army.mil/
deployment2.aspx](https://www.cs.amedd.army.mil/deployment2.aspx).

AMEDD Lessons Learned: <http://lessonslearned.amedd.army.mil>.

CALL: <http://call.army.mil/>.

Center for Pre-Deployment Medicine (CPDM): Tactical Combat Medical
Care (TCMC) Course: <https://www.us.army.mil/suite/page/312889>.

PROVIDE US YOUR INPUT

To help you access information quickly and efficiently, the Center for Army Lessons Learned (CALL) posts all publications, along with numerous other useful products, on the CALL website. The CALL website is restricted to U.S. government and allied personnel.

PROVIDE FEEDBACK OR REQUEST INFORMATION

<<http://call.army.mil>>

If you have any comments, suggestions, or requests for information (RFIs), use the following links on the CALL home page: “RFI or a CALL Product” or “Contact CALL.”

**PROVIDE OBSERVATIONS, INSIGHTS, AND LESSONS (OIL) OR
SUBMIT AN AFTER ACTION REVIEW (AAR)**

If your unit has identified lessons learned or OIL or would like to submit an AAR, please contact CALL using the following information:

Telephone: DSN 552-9569/9533; Commercial 913-684-9569/9533

Fax: DSN 552-4387; Commercial 913-684-4387

NIPR e-mail address: call.rfimanager@conus.army.mil

SIPR e-mail address: call.rfiagent@conus.army.mil

Mailing Address:

Center for Army Lessons Learned
ATTN: OCC, 10 Meade Ave., Bldg. 50
Fort Leavenworth, KS 66027-1350

TO REQUEST COPIES OF THIS PUBLICATION

If you would like copies of this publication, please submit your request at: <http://call.army.mil>. Use the “RFI or a CALL Product” link. Please fill in all the information, including your unit name and official military address. Please include building number and street for military posts.

PRODUCTS AVAILABLE “ONLINE”

CENTER FOR ARMY LESSONS LEARNED

Access and download information from CALL’s website. CALL also offers Web-based access to the CALL Archives. The CALL home page address is:

<<http://call.army.mil>>

CALL produces the following publications on a variety of subjects:

- **Combat Training Center Bulletins, Newsletters, and Trends**
- **Special Editions**
- ***News From the Front***
- **Training Techniques**
- **Handbooks**
- **Initial Impressions Reports**

You may request these publications by using the “RFI or a CALL Product” link on the CALL home page.

**COMBINED ARMS CENTER (CAC)
Additional Publications and Resources**

The CAC home page address is:

<<http://usacac.army.mil/cac2/index.asp>>

Center for Army Leadership (CAL)

CAL plans and programs leadership instruction, doctrine, and research. CAL integrates and synchronizes the Professional Military Education Systems and Civilian Education System. Find CAL products at <<http://usacac.army.mil/cac2/cal/index.asp>>.

Combat Studies Institute (CSI)

CSI is a military history think tank that produces timely and relevant military history and contemporary operational history. Find CSI products at <<http://usacac.army.mil/cac2/csi/csipubs.asp>>.

Combined Arms Doctrine Directorate (CADD)

CADD develops, writes, and updates Army doctrine at the corps and division level. Find the doctrinal publications at either the Army Publishing Directorate (APD) <<http://www.usapa.army.mil>> or the Reimer Digital Library <<http://www.adtdl.army.mil>>.

Foreign Military Studies Office (FMSO)

FMSO is a research and analysis center on Fort Leavenworth under the TRADOC G2. FMSO manages and conducts analytical programs focused on emerging and asymmetric threats, regional military and security developments, and other issues that define evolving operational environments around the world. Find FMSO products at <<http://fmso.leavenworth.army.mil/>>.

Military Review (MR)

MR is a revered journal that provides a forum for original thought and debate on the art and science of land warfare and other issues of current interest to the U.S. Army and the Department of Defense. Find MR at <<http://usacac.army.mil/cac2/militaryreview/index.asp>>.

TRADOC Intelligence Support Activity (TRISA)

TRISA is a field agency of the TRADOC G2 and a tenant organization on Fort Leavenworth. TRISA is responsible for the development of intelligence products to support the policy-making, training, combat development, models, and simulations arenas. Find TRISA Threats at <<https://dcsint-threats.leavenworth.army.mil/default.aspx>> (requires AKO password and ID).

Combined Arms Center-Capability Development Integration Directorate (CAC-CDID)

CAC-CDID is responsible for executing the capability development for a number of CAC proponent areas, such as Information Operations, Electronic Warfare, and Computer Network Operations, among others. CAC-CDID also teaches the Functional Area 30 (Information Operations) qualification course. Find CAC-CDID at <<http://usacac.army.mil/cac2/cdid/index.asp>>.

U.S. Army and Marine Corps Counterinsurgency (COIN) Center

The U.S. Army and Marine Corps COIN Center acts as an advocate and integrator for COIN programs throughout the combined, joint, and interagency arena. Find the U.S. Army/U.S. Marine Corps COIN Center at: <<http://usacac.army.mil/cac2/coin/index.asp>>.

Joint Center for International Security Force Assistance (JCISFA)

JCISFA's mission is to capture and analyze security force assistance (SFA) lessons from contemporary operations to advise combatant commands and military departments on appropriate doctrine; practices; and proven tactics, techniques, and procedures (TTP) to prepare for and conduct SFA missions efficiently. JCISFA was created to institutionalize SFA across DOD and serve as the DOD SFA Center of Excellence. Find JCISFA at <<https://jcsifa.jcs.mil/Public/Index.aspx>>.

Support CAC in the exchange of information by telling us about your successes so they may be shared and become Army successes.

<http://call.army.mil>



Center for Army Lessons Learned (CALL)

10 Meade Avenue, Building 50

Fort Leavenworth, KS 66027-1350

Combined Arms Center (CAC) • Ft. Leavenworth, KS



US Army
Combined
Arms Center

"Intellectual Center of the Army"

U.S. UNCLASSIFIED
REL NATO, GCTF, ISAF, ABCA
FOR OFFICIAL USE ONLY